



## Rehabilitation Protocol Anterior Capsular Shift (Open)

**DEREK SHIA, M.D**

9250 SW Hall Blvd., Tigard, OR 97223

503-293-0161

### PHASE I: PROTECT REPAIR

*(0 to 6 weeks after surgery)*

- Patients may shower post-op day # 5.
- Sutures will be removed by surgeon in 7 to 10 days.
- Ice or cold flow systems encouraged for the first week at a minimum: should be used 3-4 times per day.
- Sling should be in place when not performing exercises.
- Initiate exercise program 3 times per day:
  - Immediate elbow, forearm and hand range of motion out of sling
  - Pendulum exercises
  - Passive and active assistive ER at the side to 30, flexion to 130
- May start active scapular mobility exercises at 3 to 4 weeks – Must keep the shoulder musculature relaxed.
- AVOID range of motion into abduction, ER >30 degrees or ACTIVE IR

### PHASE II: PROGRESS ROM & PROTECT REPAIR

*(6 to 12 weeks after surgery)*

- May discontinue sling.
- Lifting restriction of 5 pounds with the involved extremity.
- Advance active and passive ROM:
  - ER at the side and flexion to tolerance
  - Scapular plane elevation to 130
  - IR and extension to tolerance
- Initiate gentle rotator cuff strengthening, EXCEPT IR.
- Continue scapular stabilizer strengthening.
- Avoid combined abduction and ER ROM, active or passive.
- Avoid active or resistive IR.

### PHASE III: FULL FUNCTION

*(>3 months after surgery)*

- Begin combined abduction and ER ROM and capsular mobility.
- Discontinue lifting restrictions.
- Begin subscapularis strengthening.
- Advance rotator cuff and scapular stabilizer strengthening.
- Initiate functional progression to sports specific activities at 4 months.